

Extracting Gromov-Witten invariants of a conifold from semi-stable reduction and relative GW invariants of pairs

Chien-Hao Liu and Shing-Tung Yau

(From C.-H.L. : In memory of Barbara M. Willman.)

Abstract

The study of open/closed string duality and large N duality suggests a Gromov-Witten theory for conifolds that sits on the border of both a closed Gromov-Witten theory and an open Gromov-Witten theory. The work of Jun Li on Gromov-Witten theory for a projective singular variety of the gluing form $Y_1 \cup_D Y_2$, where D is a smooth divisor on smooth Y_1 and Y_2 , suggests two methods to study Gromov-Witten invariants for a projective conifold: one by a direct generalization of his construction to the conifold singularity and the other by an appropriate semi-stable reduction of a degeneration to a conifold and then apply his results on this new degeneration to extract Gromov-Witten invariants of the original conifold. In this work we carry out the second method. Suggested by the semi-stable reduction, we associate to a conifold Y with singular locus $\{p_1, \dots\}$ a set of smooth variety-divisor pairs $(\tilde{Y}, E)$, (Y_i, D_i) , $i = 1, \dots$, and a canonical morphism $\tilde{Y} \cup_{\coprod_i D_i} \coprod_i Y_i \rightarrow Y$, where $\tilde{Y}$ is the blow-up of Y at the conifold singularities, $E \simeq \coprod_i D_i$ is the exceptional divisor, and Y_i is a smooth quadric hypersurface in $\mathbb{P}^4$ with a smooth hyperplane section $D_i \simeq \mathbb{P}^1 \times \mathbb{P}^1$, and i runs through the labels of the conifold singularities p_i . The existence of a $\mathbb{Z}/2\mathbb{Z}$ -action on the quadric hypersurface Y_i that restricts to an exchange of the two product factor $\mathbb{P}^1$'s of D_i implies that the variety $\tilde{Y} \cup_{\coprod_i D_i} \coprod_i Y_i$ from gluing these pairs is uniquely determined by Y up to isomorphisms. Jun Li's relative Gromov-Witten theory and invariants for smooth variety-divisor pairs $(Y_0, \coprod_i D_i)$ and (Y_i, D_i) and a refinement of his degeneration formula worked out here for the current situation are then employed to extract Gromov-Witten invariants of Y .

Key words: string world-sheet instanton, open/closed string duality, conifold, stable morphism, Gromov-Witten invariant, semi-stable reduction, quadric hypersurface, degeneration formula.


MSC number 2000: 14N35, 81T30.

Acknowledgements. We thank Joe Harris, Jun Li, Kefeng Liu, and Cumrun Vafa for many influences on this work via lectures, communications, and/or very profound works. C.-H.L. would like to thank in addition K.L. for serving as a sounding board to many of his raw ideas; Lisa Randall, Itay Yavin for valuable lectures; Andrew Strominger, Xiaowei Wang for discussions; Department of Mathematics of UCLA for hospitality; Rev. Campbell Willman and Ann Willman for a conversation; and Ling-Miao Chou for the tremendous moral support. The work is supported by NSF grants DMS-9803347 and DMS-0074329.

0. Introduction and outline.

Introduction: Gromov-Witten theory for conifolds in an open/closed string duality and a large N duality.

Given the 3-sphere S^3 , let X_0 be the conifold from the degeneration of $X := T^*S^3$, as a complex 3-fold, that pinches the zero-section 3-cycle S^3 of T^*S^3 and X' be the complex 3-fold from the small resolution of X_0 with exceptional locus $\simeq \mathbb{P}^1$. (X' is isomorphic to the total space of the bundle $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$ and is a non-compact Calabi-Yau 3-fold.) Gopakumar and Vafa [G-V] conjecture the following correspondence that relates the $U(N)$ or $SU(N)$ Chern-Simons gauge theory on S^3 and an A-model topological closed string theory on X' via the mechanisms indicated below:



Under this correspondence,

$U(N)$ or $SU(N)$ Chern-Simons gauge theory on S^3	$\Longleftrightarrow$	A-model topological closed string theory on X'
· 't Hooft coupling $\lambda := Ng_s$ · Wilson loop observable · large N limit of Chern-Simons gauge theory on S^3		· B-field magnitude on $\mathbb{P}^1$ · quantity in the effective theory of associated brane-probe in X' · A-model topological string theory on conifold X_0

This diagram involves a large N duality and an open/closed string duality, and an A-model topological string theory with a conifold as the target space serves as a geometric mediator and transition-point for these dualities. The dualities were tested/examined from *five* different aspects of stringy dualities: the *string world-sheet* aspect, the *target space-time* aspect, the *low dimensional effective field theory* aspect, the *brane-probe* aspect, and the *Wilson's theory space* aspect. See [G-P], [G-V], [O-V1], [O-V2], [Va], [Wi1], [Wi2] for string-theoretical details and insights, [G-R] for an introductory mathematical review and more references, and e.g. [A-M-V] and [D-F-G] for generalizations to more general non-compact Calabi-Yau 3-folds involving toric geometry.

Here a *conifold* X_0 (e.g. [B-L], [C-dlO-G-P], and [St]) is by definition the singular variety from a degeneration of Calabi-Yau 3-fold X via a deformation of complex structures

that pinches isolated smoothly-embedded 3-spheres S^3 in a smooth 3-fold X . And the partition function of an A-model topological string theory with target X_0 is supposed to compute the string world-sheet instanton numbers that can be interpreted either as a counting of holomorphic maps from (complex) curves to X_0 or from bordered Riemann surfaces to X_0 with boundary components, if non-empty, mapped to the isolated singularities of X_0 . In other words, this is a *Gromov-Witten theory for conifolds* on the mathematical side. And the open/closed string duality from string theory reveals a very distinguished feature of it as a theory on the border of *both* a closed Gromov-Witten theory on one geometry and an *open* Gromov-Witten theory on another geometry related to the previous geometry via the conifold transition.

To develop a Gromov-Witten theory for conifolds, there are two paths one may attempt to follow. The first one is to generalize the techniques in [Li1] and [Li2] directly to a conifold singularity. This is technical. The second one is to try to replace a conifold as the degenerate fiber of a smooth $\mathbb{A}^1$ -family $W/\mathbb{A}^1$ by the degenerate fiber of a semi-stable reduction of $W/\mathbb{A}^1$ and see if one can reduce the problem to the case already dealt with in [Li1] and [Li2] and use it to extract the Gromov-Witten invariants of the conifold. Surely, for general singularities one will not expect the second path would immediately work either since one still misses the understanding of Gromov-Witten theory for a pair (Y, D) , where Y is smooth and D is a divisor on Y with simple normal crossing singularities, and a related degeneration/gluing formula for gluing Gromov-Witten invariants from a collection of such pairs. As if given by God, it turns out that such potentially existing difficulties along the second path are not really there for a conifold singularity, (Sec. 1). This accidental simplicity for conifolds together with similar discussions and results in [L-Y] that refine Jun Li's degeneration formula to one for curve classes in $H_2(\cdot; \mathbb{Z})$ or $A_1(\cdot)$, (Sec. 2), enable us to extract some Gromov-Witten invariants of conifolds, (Sec. 3). We explain the details of this second path in this work.

Convention. This work is parallel to [L-Y] and follows the notations and the terminology of [Li1], [Li2], and [L-Y] closely, except where confusions may occur. Other notations follow [Hart], [Fu], [De], and [K-M]. All schemes are over $\mathbb{C}$ and all points are referred to closed points. All conifolds are assumed to be projective. Though appearing a few times in the Introduction for easy match with literatures in string theory, the Calabi-Yau condition is not relevant in this work but will be relevant in an application.

Preliminary. Readers are referred to [Li1: Sec. 0], [Li2: Sec. 0], and [L-Y: Sec. 1, Sec. 2] for definitions and an outline of Jun Li's work that are needed for the current work.

Outline.

1. A semi-stable reduction of a conifold degeneration.
2. A degeneration formula of Gromov-Witten invariants with respect to a curve class.
3. Extracting Gromov-Witten invariants of a conifold from pairs.

1 A semi-stable reduction of a conifold degeneration.

Semi-stable reduction and the associated smooth pairs of a conifold.

Let $\pi : W \rightarrow \mathbb{A}^1$ be a flat family of (complex) 3-dimensional projective varieties with smooth general fibers W_t , $t \neq \mathbf{0}$, and smooth total space W such that W_t degenerates to a conifold $W_0 = Y$ over $\mathbf{0} \in \mathbb{A}^1$. For simplicity of notations and presentations, we assume that Y has only one conifold singularity. An example of such family can be obtained from a degeneration of quintic 3-folds in $\mathbb{P}^4$, cf. [C-dlO-G-P]. In a local analytic germ, the degeneration to a conifold singularity is modeled on the morphism

$$\begin{array}{ccc} \text{Spec } \mathbb{C}[[x, y, z, w]] & \longrightarrow & \text{Spec } \mathbb{C}[[t]] \\ xy - zw & \longleftarrow & t. \end{array}$$

This gives a 3-dimensional isolated hypersurface singularity of multiplicity 2 at the fiber over $\mathbf{0} := (t)$. The family $W/\mathbb{A}^1$ can be semi-stabilized by the following sequence of blow-ups and finite base change obtained from a straightforward computation:

$$\begin{array}{ccccccc} W^{(3)} & \xrightarrow{\varphi_3} & W^{(2)} & \xrightarrow{\varphi_2} & W^{(1)} & \xrightarrow{\varphi_1} & W^{(0)} := W \\ \pi_3 \downarrow & & \pi_2 \downarrow & & \downarrow \pi_1 & & \downarrow \pi_0 = \pi \\ \mathbb{A}^1 & = & \mathbb{A}^1 & \xrightarrow{\alpha} & \mathbb{A}^1 & = & \mathbb{A}^1 \end{array},$$

where

- $\varphi_1 : W^{(1)} \rightarrow W^{(0)}$ is the blow-up of W at the conifold singularity of W_0 ;
- recall that W is smooth and hence the exceptional locus of φ_1 is a $\mathbb{P}^3$; the degenerate fiber $\pi_1^{-1}(\mathbf{0})$ of π_1 contains a nonreduced irreducible component of multiplicity 2 supported on this $\mathbb{P}^3$; as divisors in $W^{(1)}$, $\pi_1^{-1}(\mathbf{0}) = \tilde{Y} + 2\mathbb{P}^3$, where $\tilde{Y}$ is the resolution of Y by a blow-up at the conifold singularity and $\tilde{Y} \cap \mathbb{P}^3$ is a smooth quadric surface in $\mathbb{P}^3$;
- $\alpha : (\mathbb{A}^1, \mathbf{0}) \rightarrow (\mathbb{A}^1, \mathbf{0})$ is a finite morphism of degree 2 branched over $\mathbf{0}$, φ_2 and π_2 on $W^{(2)}$ are from the fibered-product of α and π_1 ;
- $W^{(2)}$ is now a singular scheme whose singularities are modelled on the Whitney's umbrella; the singular locus $\text{Sing}(W^{(2)})$ of $W^{(2)}$ with the reduced subscheme structure has multiplicity 2, lies over $\mathbf{0}$, and is isomorphic to $\mathbb{P}^3$;
- normalization/blow-up of $W^{(2)}$ along $\text{Sing}(W^{(2)}) \simeq \mathbb{P}^3$ gives $W^{(3)}$ which is smooth with $\pi_3^{-1}(\mathbf{0}) = Y_0 \cup Y_1$, where $Y_0 := \tilde{Y}$ and Y_1 is naturally realized as a double cover of $\mathbb{P}^3$ branched over a smooth quadric surface in $\mathbb{P}^3$.

In local germs or local formal schemes, the morphisms in the diagram above are given by

$$\begin{array}{ccc}
\begin{array}{c}
\text{Spec } \mathbb{C}[[x_1, y_1, z_1, w_1]] \\
w_1^2(x_1 y_1 - z_1) \quad x_1 w_1; y_1 w_1; z_1 w_1; w_1 \\
\pi_1^\# \uparrow \quad \pi_1 \downarrow \\
t_1 \quad \text{Spec } \mathbb{C}[[t_1]]
\end{array}
&
\begin{array}{c}
\varphi_1 \rightarrow \\
\varphi_1^\# \leftarrow \\
\leftarrow \quad \rightarrow
\end{array}
&
\begin{array}{c}
\text{Spec } \mathbb{C}[[x_0, y_0, z_0, w_0]] \\
x_0 y_0 - z_0 w_0 \\
\downarrow \pi_0 \quad \uparrow \pi_0^\# \\
t_0 \quad \text{Spec } \mathbb{C}[[t_0]]
\end{array}
, \\
\\
\begin{array}{c}
\text{Spec } \mathbb{C}[[x_1, y_1, z_1, w_1, t_2]]/(t_2^2 - w_1^2(x_1 y_1 - z_1)) \\
t_2 \quad x_1; y_1; z_1; w_1 \\
\pi_2^\# \uparrow \quad \pi_2 \downarrow \\
t_2 \quad \text{Spec } \mathbb{C}[[t_2]]
\end{array}
&
\begin{array}{c}
\varphi_2 \rightarrow \\
\varphi_2^\# \leftarrow \\
\leftarrow \quad \rightarrow
\end{array}
&
\begin{array}{c}
\text{Spec } \mathbb{C}[[x_1, y_1, z_1, w_1]] \\
w_1^2(x_1 y_1 - z_1) \\
\downarrow \pi_1 \quad \uparrow \pi_1^\# \\
t_1 \quad \text{Spec } \mathbb{C}[[t_1]]
\end{array}
;
\end{array}$$

consider the ring isomorphism generated by $x_1 \mapsto x_2$, $y_1 \mapsto y_2$, $z_1 \mapsto x_2 y_2 - z_2$, $w_1 \mapsto w_2$, and $t_2 \mapsto t_2$ and rewrite

$$\text{Spec } \mathbb{C}[[x_1, y_1, z_1, w_1, t_2]]/(t_2^2 - w_1^2(x_1 y_1 - z_1)) \quad \text{as} \quad \text{Spec } \mathbb{C}[[x_2, y_2, z_2, w_2, t_2]]/(t_2^2 - w_2^2 z_2),$$

which reveals the Whitney umbrella transverse surface singularity of $W^{(2)}$; and then

$$\begin{array}{ccc}
\begin{array}{c}
\text{Spec } \mathbb{C}[[x_3, y_3, z_3, w_3]] \\
z_3 w_3 \quad x_3; y_3; z_3^2; w_3; z_3 w_3 \\
\pi_3^\# \uparrow \quad \pi_3 \downarrow \\
t_3 \quad \text{Spec } \mathbb{C}[[t_3]]
\end{array}
&
\begin{array}{c}
\varphi_3 \rightarrow \\
\varphi_3^\# \leftarrow \\
\leftarrow \quad \rightarrow
\end{array}
&
\begin{array}{c}
\text{Spec } \mathbb{C}[[x_2, y_2, z_2, w_2, t_2]]/(t_2^2 - w_2^2 z_2) \\
x_2; y_2; z_2; w_2; t_2 \\
\downarrow \pi_2 \quad \uparrow \pi_2^\# \\
t_2 \quad \text{Spec } \mathbb{C}[[t_2]]
\end{array}
.
\end{array}$$

(Cf. FIGURE 1-1.)




FIGURE 1-1. A semi-stable reduction of a conifold degeneration.
The degenerate fiber over $0 \in \mathbb{A}^1$ in each family is indicated.

Definition 1.1 [canonical semi-stable reduction]. We shall call $W^{(3)}/\mathbb{A}^1$ obtained above the *canonical semi-stable reduction* of the conifold degeneration $W/\mathbb{A}^1$.

After this semi-stable reduction of the conifold degeneration $\pi : W \rightarrow \mathbb{A}^1$, one obtains a new family $\pi_3 : W^{(3)} \rightarrow \mathbb{A}^1$ of 3-folds with $W^{(3)}$ smooth, $W_t^{(3)} = W_{t^2}$ for $t \neq 0$, and $W_0^{(3)} := \pi_3^{-1}(0) = Y_0 \cup_D Y_1$ is a variety with simple normal crossing singularity from gluing smooth Y_0 and Y_1 along the isomorphic smooth divisor D . This happens to be exactly the type of degenerations whose Gromov-Witten theory are studied in [Li1] and [Li2].

Definition 1.2 [associated canonical pairs to conifold]. We shall call the set of pairs, (Y_0, D) and (Y_1, D) , the *set of canonical smooth pairs associated to the conifold Y* .

By construction, there is a canonical morphism $p = p_0 \cup_D p_1 : Y_0 \cup_D Y_1 \rightarrow Y$, where $p_0 : Y_0 \rightarrow Y$ is the resolution of Y whose exception locus is the smooth divisor D and $p_1 : Y_1 \rightarrow Y$ pinches Y_1 to the conifold singularity of Y .

Special properties of Y_1 .

The 3-fold Y_1 is naturally realized as a quadric hypersurface in $\mathbb{P}^4$ with $D \simeq \mathbb{P}^1 \times \mathbb{P}^1$ realized as the intersection of Y_1 with a hyperplane H in $\mathbb{P}^4$. It follows from the Bruhat cellular decomposition of a smooth quadric hypersurface in a projective space and the Lefschetz hyperplane theorem that the only non-vanishing Chow groups and (singular) homology groups for Y_1 coincide and are given by

$$A_k(Y_1) \simeq H_{2k}(Y_1; \mathbb{Z}) \simeq \mathbb{Z}, \quad \text{for } k = 0, 1, 2, 3.$$

In particular, $A_1(Y_1) \simeq H_2(Y_1; \mathbb{Z})$ is generated by a complex line γ_2 from either of the rulings of D , $A_2(Y_1) \simeq H_4(Y_1; \mathbb{Z})$ is generated by the complex surface $\gamma_4 := D$, and $\gamma_2 \cap \gamma_4 = +1$ on Y_1 . (See [Fu], [G-H], [Harr], and [P-S] for details.)

Fix an embedding $D = Y_1 \cap H \subset Y_1 \hookrightarrow \mathbb{P}^4$ and an isomorphism $D \simeq \mathbb{P}^1 \times \mathbb{P}^1$ from the rulings of D . Up to a $PGL(5, \mathbb{C})$ -action on $\mathbb{P}^4$, one can write this explicitly in terms of homogeneous coordinates $[u : x : y : z : w]$ of $\mathbb{P}^4$ as

$$D = V(u, u^2 + xy - zw) \subset Y_1 = V(u^2 + xy - zw) \subset \mathbb{P}^4 \quad \text{with} \quad H = V(u).$$

The $\mathbb{Z}/2\mathbb{Z}$ -action on D that interchanges the two $\mathbb{P}^1$ -factors extends to a linear $\mathbb{Z}/2\mathbb{Z}$ -action on $\mathbb{P}^4$ that leaves Y_1 invariant. Explicitly, $D \simeq \mathbb{P}^1 \times \mathbb{P}^1$ via $\{xy - zw = 0\} \rightarrow ([x : z], [x : w])$ and one can choose this $\mathbb{Z}/2\mathbb{Z}$ -action to be the exchange $z \leftrightarrow w$. It follows that:

Lemma 1.3 [uniqueness of gluing]. *Any variety from gluing the smooth pairs, (Y_0, D) and (Y_1, D) , associated to Y via an automorphism of D is isomorphic to each other.*

The monodromy of $\pi : W \rightarrow \mathbb{A}^1$.

To be able to refine Jun Li's degeneration formula from one with respect to a π_3 -ample line bundle on $W^{(3)}/\mathbb{A}^1$ to one with respect to a curve class of a general fiber W_t of $W/\mathbb{A}^1$, the

monodromy around the degenerate fiber $W_0 = Y$ of $W/\mathbb{A}^1$ is required to be well-behaved. Let us thus take a look at this. In this part, we will leave the algebro-geometric category over $\mathbb{C}$ and enter the topological category over $\mathbb{R}$. In particular, ‘ $\simeq$ ’ means ‘topologically homeomorphic’.

Topologically, the degeneration $W/\mathbb{A}^1$ pinches a real 3-cycle realized as a smoothly embedded 3-sphere S^3 in W_t , $t \neq \mathbf{0}$, to the conifold singularity of $W_0 = Y$. Over a loop $S^1 \hookrightarrow \mathbb{A}^1$ around $\mathbf{0} \in \mathbb{A}^1$ (here we will think of $\mathbb{A}^1$ as the usual complex plane $\mathbb{C}$ with the analytic topology and S^1 is an oriented circle therein), $\pi^{-1}(S^1)$ is the mapping torus associated to a smooth automorphism $\phi : W_t \xrightarrow{\sim} W_t$. ϕ can be homotoped to an automorphism, still denoted by ϕ , on W_t that is not the identity map only in a small tubular neighborhood $N_\varepsilon(S^3/W_t) \simeq S^3 \times B^3$ of S^3 in W_t , where B^3 is the unit (real) 3-ball and there is an orientation on (S^3, B^3) that is compatible with the complex structure on $N_\varepsilon(S^3/W_t)$. Indeed, ϕ is a *generalized Dehn twist* along S^3 and, up to homotopy, can be described explicitly as follows. Recall that S^3 is the underlying topology of the Lie group $SU(2)$. Let e be its identity element and fix a degree 1 map $\phi_e : (B^3, \partial B^3) \rightarrow (S^3, e)$ that is a homeomorphism from $B^3 - \partial B^3$ to $S^3 - \{e\}$. Define $\phi : N_\varepsilon(S^3/W_t) \rightarrow N_\varepsilon(S^3/W_t)$ by

$$\begin{aligned} \phi : S^3 \times B^3 &\longrightarrow S^3 \times B^3 \\ (g, x) &\longmapsto (g \cdot \phi_e(x), x) \quad , \end{aligned}$$

where the $\cdot$ in the definition is the group multiplication of $SU(2)$. By construction, ϕ restricts to the identity map on $\partial(N_\varepsilon(S^3/W_t))$ and extends by the identity map to the automorphism ϕ on W_t .

Since $H_i(S^3 \times B^3, \partial(S^3 \times B^3); \mathbb{Z}) = 0$ for $i = 0, 1, 2$, the monodromy of ϕ on $H_i(W_t; \mathbb{Z})$ is the identity map for $i = 0, 1, 2$. (The monodromy on $H_3(W_t; \mathbb{Z})$ is the much studied Picard-Lefschetz operation $\gamma \mapsto \gamma + \langle \gamma, [S^3] \rangle [S^3]$, where $\langle \cdot, \cdot \rangle$ is the intersection pairing on $H_3(W_t; \mathbb{Z})$. But we do not need this for this work.) Since the conifold $W_0 = Y$ is topologically the quotient space of W_t , $t \neq \mathbf{0}$, by identifying the vanishing 3-cycle S^3 to a point, Y is homotopically equivalent to $W_t \cup_{S^3 \simeq \partial B^4} B^4$, where $t \neq \mathbf{0}$ and the 4-ball B^4 is attached to W_t along an isomorphism of ∂B^4 to the vanishing 3-cycle S^3 . This implies that $H_i(W_0; \mathbb{Z}) \simeq H_i(W_t; \mathbb{Z})$ canonically for $i = 0, 1, 2$. In particular, $R^\bullet \pi_* \mathbb{Z}_W$ has a summand that is the constant sheaf on $\mathbb{A}^1$ associated to the group $H_2(W_t; \mathbb{Z}) = H_2(Y; \mathbb{Z})$.

To summarize,

Lemma 1.4 [canonical identification and trivial monodromy on H_2]. *The monodromy of $W/\mathbb{A}^1$ around the conifold $W_0 = Y$ is trivial on $H_2(W_t; \mathbb{Z})$. Each $H_2(W_t; \mathbb{Z})$, $t \neq \mathbf{0}$, is canonically isomorphic to $H_2(W_0; \mathbb{Z}) = H_2(Y; \mathbb{Z})$.*

We now resume to proceed in the domain of algebraic geometry.

2 A degeneration formula of Gromov-Witten invariants with respect to a curve class.

Once the semi-stable degeneration $W^{(3)}/\mathbb{A}^1$ of the conifold degeneration $W/\mathbb{A}^1$ is understood, the second ingredient toward a Gromov-Witten theory for Y via extracting information from gluing pairs (Y_0, D) and (Y_1, D) is the degeneration/gluing formula of Gromov-Witten invariants for the family $W^{(3)}/\mathbb{A}^1$ that is intrinsic to curve classes on Y (note: *not* on $W_0^{(3)}$). The goal of this section is to derive such a formula from [Li2]. The reasoning and the discussion of this section are parallel to [L-Y: Sec. 2 and Sec. 3]. Readers are referred to [L-Y] for more explanations in a simpler situation and [Li1] and [Li2] for the thorough related details.

A refinement of Jun Li's degeneration formula.

Recall first the definitions of the following: *admissible weighted graph* ([Li1: Definition 4.6], [L-Y: Definition 1.1]; *admissible triple* ([Li1: Definition 4.11], [L-Y: Definition 1.2]), the integer triple $|\eta|$ for an admissible triple $\eta = (\Gamma_0, \Gamma_1, I)$, and Jun Li's *degeneration formula* ([Li2: Sec. 0, Theorem 3.15, and Corollary 3.16]).

Fix a π_3 -ample line bundle $\mathcal{L}$ on $W^{(3)}/\mathbb{A}^1$ and an $\mathcal{L}$ -degree d , then the stack of stable morphisms from prestable curves of genus g , n marked points, into fibers of the universal family of expanded degenerations associated to $W^{(3)}/\mathbb{A}^1$ of $\mathcal{L}$ -degree d is a Deligne-Mumford stack $\mathfrak{M}(\mathfrak{W}^{(3)}, (g, n; d))$ with a perfect obstruction theory. By construction, $\mathfrak{M}(\mathfrak{W}^{(3)}, (g, n; d))$ fibers naturally over $\mathbb{A}^1$ with the fiber over $\mathbf{0}$ denoted by $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; d))$. The perfect obstruction theory on $\mathfrak{M}(\mathfrak{W}^{(3)}, (g, n; d))$ restricts to a perfect obstruction theory on $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; d))$. The Gromov-Witten theory and invariants of the singular variety $Y_0 \cup_D Y_1$ from gluing is defined via the virtual fundamental class $[\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; d))]^{virt}$ on $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; d))$. On the other hand, for each of the smooth variety-divisor pairs, (Y_i, D) , Jun Li constructed a general theory of relative Gromov-Witten invariants, which consists of the moduli stack $\mathfrak{M}(\mathfrak{Y}_i^{rel}, \Gamma_i)$ of stable morphisms of topological type Γ_i from prestable curves to the fibers of the universal family of the stack $\mathfrak{Y}_i^{rel}$ of expanded relative pairs associated to (Y_i, D) and a perfect obstruction theory thereon. For each admissible triple η with $|\eta| = (g, n; d)$, there is a morphism, (the r below is the total root weight of Γ_0 in η),

$$\Phi_\eta : \mathfrak{M}(\mathfrak{Y}_0^{rel}, \Gamma_0) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_1^{rel}, \Gamma_1) \longrightarrow \mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; d))$$

that is finite étale to its image $\mathfrak{M}(\mathfrak{Y}_0^{rel} \sqcup \mathfrak{Y}_1^{rel}, \eta)$ and has degree a combinatorial factor $|\text{Eq}(\eta)|$ from η , cf. [Li1: Sec. 4]. Up to the difference from a nonreduced structure that has to be taken care ([Li2: Sec. 4.4]), $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; d))$ is the union of the substacks $\mathfrak{M}(\mathfrak{Y}_0^{rel} \sqcup \mathfrak{Y}_1^{rel}, \eta)$ in $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; d))$. Going from $\mathfrak{M}(\mathfrak{W}_t^{(3)}, (g, n; d))$, $t \neq \mathbf{0}$, which gives the usual Gromov-Witten theory of the smooth projective W_t to $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; d))$, which gives the Gromov-Witten theory on $Y_0 \cup_D Y_1$, and recast it to a form from gluing relative

Gromov-Witten theory of pairs, (Y_0, D) and (Y_1, D) , gives a degeneration formula that relates Gromov-Witten invariants of W_t to a combination of relative Gromov-Witten invariants of (Y_0, D) and (Y_1, D) in the degenerate fiber $W_0^{(3)}$. See [Li1] and [Li2] for the complete technical details.

Since the monodromy on $H_2(W_t; \mathbb{Z})$ around $\mathbf{0} \in \mathbb{A}^1$ is trivial, so is the monodromy on $H_2(W_t^{(3)}; \mathbb{Z})$ around $\mathbf{0}$. Recall also the canonical isomorphism $H_2(W_t; \mathbb{Z}) \simeq H_2(Y; \mathbb{Z})$. Thus, a $\beta' \in H_2(Y; \mathbb{Z})$ determines a unique class in $H_2(W_t; \mathbb{Z})$, $t \neq \mathbf{0}$, and hence a unique class, still denoted by β' , in $H_2(W_t^{(3)}; \mathbb{Z})$, $t \neq \mathbf{0}$. Define the $\mathcal{L}$ -degree $\mathcal{L} \cdot \beta'$ of $\beta' \in H_2(Y; \mathbb{Z})$ via this identification. Let d be the $\mathcal{L}$ -degree for a given curve class $\beta \in H_2(Y; \mathbb{Z})$ and define $C_{(\mathcal{L}, d)}(Y) := \{ \beta' \in H_2(Y; \mathbb{Z}) : \mathcal{L} \cdot \beta' = d \}$. Then

$$\mathfrak{M}(\mathfrak{W}^{(3)}, (g, n; d)) = \coprod_{\beta' \in C_{(\mathcal{L}, d)}(Y)} \mathfrak{M}(\mathfrak{W}^{(3)}, (g, n; \beta')),$$

where $\mathfrak{M}(\mathfrak{W}^{(3)}, (g, n; \beta'))$ is the stack of stable morphisms of $\mathcal{L}$ -degree d from prestable curves of genus g with n marked points to the fibers of the universal family of the stack $\mathfrak{W}^{(3)}$ of expanded degenerations associated to $W^{(3)}/\mathbb{A}^1$ such that after the post-composition with the morphisms

$$\mathfrak{W}^{(3)}/\mathbb{A}^1 \longrightarrow W^{(3)}/\mathbb{A}^1 \xrightarrow{\varphi_1 \circ \varphi_2 \circ \varphi_3} W/\mathbb{A}^1,$$

the images of the stable morphisms lie in the curve class $\beta' \in H_2(Y; \mathbb{Z})$. From its definition, $\mathfrak{M}(\mathfrak{W}^{(3)}, (g, n; \beta'))$ depends not just on (g, n, β') but also on $\mathcal{L}$. (Indeed, since Y_1 is mapped to the conifold singularity of Y , the choice of curve classes from $H_2(Y_0 \cup_D Y_1; \mathbb{Z})$ does depend on $\mathcal{L}$ in general.)

Recall the $\mathcal{L}$ -dependent set $\Omega_{(g, k; d)}$ of admissible triples η such that $|\eta| = (g, k; d)$ and the quotient set $\overline{\Omega}_{(g, k; d)}$ defined in [Li2: Sec. 0]. Recall also the morphism $p = p_0 \cup_D p_1 : Y_0 \cup_D Y_1 \rightarrow Y$. For an admissible weighted graph Γ for a relative pair, let $b(\Gamma) := \sum_{v \in V(\Gamma)} b(v)$. Define the β -compatible subset of $\Omega_{(g, n; d)}$ by

$$\Omega_{(g, k; \beta)}^{\mathcal{L}} := \{ \eta = (\Gamma_0, \Gamma_1, I) \in \Omega_{(g, k; d)} \mid p_{0*} b(\Gamma_0) = \beta \}.$$

Same discussions as in [L-Y: Sec. 2] imply the existence of a perfect obstruction theory on the moduli stack $\mathfrak{M}(\mathfrak{W}^{(3)}, (g, n; \beta))$ over $\mathbb{A}^1$ and its fiber $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; \beta))$ over $\mathbf{0}$, inherited from those constructed in [Li2]. This gives a well-defined Gromov-Witten theory and Gromov-Witten invariants on $Y_0 \cup_D Y_1$ associated to $\beta \in H_2(Y; \mathbb{Z})$, Jun Li's degeneration formula [Li2] implies then the following degeneration formula:

Lemma 2.1 [Jun Li's degeneration formula]. (Cf. [L-Y: Corollary 2.2] and explanations of notations there and in [Li1] and [Li2].) *Let $\alpha \in H_c^0(R^* \pi_{3*} \mathbb{Q}_{W^{(3)}})^{\times n}$, whose restriction to $W_t^{(3)}$ will be denoted by $\alpha(t)$, and $\zeta \in A_*(\mathfrak{M}_{g, n})$. Denote by $\Psi_{(g, n; \beta)}^{W_t^{(3)}}(\alpha(t), \zeta)$ the usual Gromov-Witten invariant of $W_t^{(3)}$ associated to these data. For $\eta \in \overline{\Omega}_{(g, n; \beta)}^{\mathcal{L}}$,*

assume that $G_\eta^*(\zeta) = \sum_{j \in K_\eta} \zeta_{\eta,0,j} \boxtimes \zeta_{\eta,1,j}$, where $G_\eta : \mathfrak{M}_{\Gamma_0^\circ} \times \mathfrak{M}_{\Gamma_1^\circ} \rightarrow \mathfrak{M}_{g,n}$ is the natural morphism between the related moduli stack of nodal curves. Then

$$\Psi_{(g,n;\beta)}^{W_t^{(3)}}(\alpha(t), \zeta) = \sum_{\eta \in \overline{\Omega}_{(g,n;\beta)}^\mathcal{L}} \frac{\mathbf{m}(\eta)}{|\text{Eq}(\eta)|} \sum_{j \in K_\eta} \left[\Psi_{\Gamma_0}^{Y_0^{rel}}(j_0^* \alpha(0), \zeta_{\eta,0,j}) \bullet \Psi_{\Gamma_1}^{Y_1^{rel}}(j_1^* \alpha(0), \zeta_{\eta,1,j}) \right]_0,$$

where $j_i : Y_i \rightarrow W_0^{(3)}$ is the inclusion map,

$$\Psi_{\Gamma_i}^{Y_i^{rel}}(j_i^* \alpha(0), \zeta_{\eta,i,j}) = \mathbf{q}_i * \left(ev^*(j_i^* \alpha(0)) \cdot \pi_{\Gamma_i}^*(\zeta_{\eta,i,j}) \cdot [\mathfrak{M}(\mathfrak{Y}_i^{rel}, \Gamma_i)]^{virt} \right) \in H_*(D^r), \quad i = 0, 1.$$

In cycle form,

$$[\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; \beta))]^{virt} = \sum_{\eta \in \overline{\Omega}_{(g,n;\beta)}^\mathcal{L}} \frac{\mathbf{m}(\eta)}{|\text{Eq}(\eta)|} \Phi_{\eta^*} \Delta^! \left([\mathfrak{M}(\mathfrak{Y}_0^{rel}, \Gamma_0)]^{virt} \times [\mathfrak{M}(\mathfrak{Y}_1^{rel}, \Gamma_1)]^{virt} \right),$$

where $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; \beta))$ is the fiber of $\mathfrak{M}(\mathfrak{W}^{(3)}, (g, n; \beta))$ over $\mathbf{0} \in \mathbb{A}^1$, $\Delta^!$ is the Gysin map associated to the diagonal map $\Delta : D^r \rightarrow D^r \times D^r$ for the relevant D^r in each summand.

We want to make things as intrinsic to Y as possible so that we can appropriately combine the Gromov-Witten invariants of $Y_0 \cup_D Y_1$ defined from [Li2] to a quantity that is justifiable to be called a Gromov-Witten invariant of Y . In particular, we want to remove the possible $\mathcal{L}$ -dependence (so far on *both* sides of the equation) in the above gluing formula. Let us now deal with this issue.

The $\mathcal{L}$ -(in)dependence of $\Omega_{(g,k;\beta)}^\mathcal{L}$ and $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; \beta))$.

Recall from Sec. 1 the intermediate families over $\mathbb{A}^1$ that occur in the semi-stable reduction of $W/\mathbb{A}^1$:

$$\begin{array}{ccccccc} W^{(3)} & \xrightarrow{\varphi_3} & W^{(2)} & \xrightarrow{\varphi_2} & W^{(1)} & \xrightarrow{\varphi_1} & W^{(0)} := W \\ \pi_3 \downarrow & & \pi_2 \downarrow & & \downarrow \pi_1 & & \downarrow \pi_0 = \pi \\ \mathbb{A}^1 & = & \mathbb{A}^1 & \xrightarrow{\alpha} & \mathbb{A}^1 & = & \mathbb{A}^1 \end{array}.$$

Denote the exceptional divisor of φ_1 on $W^{(1)}$ by $E_1 (\simeq \mathbb{P}^3)$ and recall that the exceptional divisor of φ_3 on $W^{(3)}$ is Y_1 . Recall [C-H] or [L-Y: Remark 3.1]. Let $\mathcal{L}_0$ be a sufficiently very ample line bundle on $W^{(0)} = W$. Then $\mathcal{L}_1 := (\varphi_1^* \mathcal{L}_0)(-E_1)$ is very ample on $W^{(1)}$. The pull-back $\mathcal{L}_2 := \varphi_2^* \mathcal{L}_1$ on $W^{(2)}$ is π_2 -ample. Thus there is an open subset U of $\mathbb{A}^1$, containing $\mathbf{0}$, such that $\mathcal{L}_2$ is ample on $\pi_2^{-1}(U)$. By removing finitely many fibers of the families/ $\mathbb{A}^1$ in the above diagram and with an abuse of notation that we denote $\pi_2^{-1}(U)$ over U still by $W^{(2)}/\mathbb{A}^1$ (and since it is only a neighborhood of Y in W that matters), we will say that $\mathcal{L}_2$ is ample on $W^{(2)}$. By taking $k \gg 0$, $\mathcal{L}_2^{\otimes k}$ becomes sufficiently very ample on $W^{(2)}$ and its pull-back with a twist $\mathcal{L} := (\varphi_3^* \mathcal{L}_2^{\otimes k})(-Y_1)$ becomes very ample on $W^{(3)}$.

Lemma 2.2 [$\mathcal{L}$ -independence of $\Omega_{(g,n;\beta)}^{\mathcal{L}}$]. *Let $\mathcal{L}$ be a very ample line bundle on $W^{(3)}$ as constructed above and d be the $\mathcal{L}$ -degree of $\beta \in H_2(Y; \mathbb{Z})$. Then $\Omega_{(g,n;\beta)}^{\mathcal{L}}$ that appears in the degeneration formula in Lemma 2.1 is independent of $\mathcal{L}$.*

Proof. The proof follows the same reasoning as in the proof of [L-Y: Lemma 3.2]. Recall the blow-up resolution $p_0 : Y_0 \rightarrow Y$ with exceptional divisor identical to $D (= Y_0 \cap Y_1)$ and p_1 that sends Y_1 to the conifold singularity of Y . It follows from a careful chasing through the sequence of pull-backs and twists in the construction of $\mathcal{L}$ that

$$\mathcal{L}^{(0)} := \mathcal{L}|_{Y_0} \simeq (p_0^* \mathcal{L}_0|_Y)^{\otimes k} (-(k+1)D) \quad \text{and} \quad \mathcal{L}^{(1)} := \mathcal{L}|_{Y_1} \simeq \mathcal{O}_{Y_1}(-(k+1)D).$$

Recall the generator γ_2 of $H_2(Y_1; \mathbb{Z})$. Let $\gamma_{2,1}$ and $\gamma_{2,2}$ be the curve classes on Y_0 from the two rulings of D . Then they generate the semi-group of curve classes in the relative Mori cone $NE(p_0)$. Observe that $D \cdot \gamma_{2,i} = -1$ on Y_0 for $i = 1, 2$ while $D \cdot \gamma_2 = +1$ on Y_1 . Let $NE(Y)_\beta$ be the set of all the curve classes in $NE(Y)_\mathbb{Z}$ that represent β in $H_2(Y; \mathbb{Z})$. Then, for each $\gamma \in NE(Y)_\beta$, $p_{0*}^{-1}(\gamma)$ in $NE(Y_0)_\mathbb{Z}$ is of the form $\tilde{\gamma} + \mathbb{Z}_{\geq 0} \gamma_{2,1} + \mathbb{Z}_{\geq 0} \gamma_{2,2}$ for a unique $\tilde{\gamma} \in NE(Y_0)_\mathbb{Z}$ determined by γ .

Now let $\eta = (\Gamma_0, \Gamma_1, I) \in \Omega_{(g,n;\beta)}^{\mathcal{L}}$. Then the pairs $(b(\Gamma_0), b(\Gamma_1))$ are characterized by the conditions:

$$p_{0*} b(\Gamma_0) = \beta \quad \text{and} \quad \mathcal{L}^{(0)} \cdot b(\Gamma_0) + \mathcal{L}^{(1)} \cdot b(\Gamma_1) = \mathcal{L} \cdot \beta = d.$$

Solving it explicitly as in [L-Y: proof of Lemma 3.2], one concludes that

$$\Omega_{(g,n;\beta)}^{\mathcal{L}} = \coprod_{\gamma \in NE(Y)_\beta} \left\{ \begin{array}{l} \eta = (\Gamma_0, \Gamma_1, I) \\ \text{admissible} \\ \text{triple} \\ \text{for } Y_0 \cup_D Y_1 \end{array} \left| \begin{array}{l} \bullet \ b(\Gamma_0) = \tilde{\gamma} + l_{0,1} \gamma_{2,1} + l_{0,2} \gamma_{2,2}, \ b(\Gamma_1) = l_1 \gamma_2, \\ \quad (l_{0,1} + l_{0,2}) + l_1 = D \cdot \tilde{\gamma}, \ l_{0,1}, l_{0,2}, l_1 \in \mathbb{Z}_{\geq 0}; \\ \bullet \ g(\eta) = g, \ k_1 + k_2 = n; \\ \bullet \ \sum_i \mu_{0,i} = l_1; \\ \bullet \ I \subset \{1, \dots, n\}, \ |I| = k_1. \end{array} \right. \right\}$$

This set is indeed independent of $\mathcal{L}$. □

We shall denote the $\mathcal{L}$ -independent set of admissible triples worked out explicitly in the end of the proof above by $\Omega_{(g,n;\beta)}$.

The proof of the above lemma implies also that, with this choice of $\mathcal{L}$ on $W^{(3)}/\mathbb{A}^1$, the potentially $\mathcal{L}$ -dependent stack $\mathfrak{M}(\mathfrak{W}^{(3)}, (g, n; \beta))$ can be re-defined without referring to $\mathcal{L}$ at all and, hence, depends only on W and $(g, k; \beta)$. Its fiber $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; \beta))$ over $\mathbf{0} \in \mathbb{A}^1$ is thus also $\mathcal{L}$ -independent. Adding this $\mathcal{L}$ -independence into the statements in Lemma 2.1, we obtain:

Theorem 2.3 [intrinsic to $W/\mathbb{A}^1$ and β]. (Cf. [L-Y: Theorem 3.3].) *The degeneration/gluing formulas, one in the numerical form and the other in the equivalent cycle*

form, in Lemma 2.1 are independent of $\mathcal{L}$ and are intrinsic to W and $(g, n; \beta)$ when $\mathcal{L}$ is chosen as above: (the cycle form omitted)

$$\Psi_{(g, n; \beta)}^{W_t^{(3)}}(\alpha(t), \zeta) = \sum_{\eta \in \overline{\Omega}_{(g, n; \beta)}} \frac{\mathbf{m}(\eta)}{|\text{Eq}(\eta)|} \sum_{j \in K_\eta} \left[\Psi_{\Gamma_0}^{Y_0^{rel}}(j_0^* \alpha(0), \zeta_{\eta, 0, j}) \bullet \Psi_{\Gamma_1}^{Y_1^{rel}}(j_1^* \alpha(0), \zeta_{\eta, 1, j}) \right]_0.$$

Remark 2.4 [W -dependence of $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; \beta))$]. From the details in [Li2: Sec. 4.4], the dependence of the stack $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; \beta))$ on the neighborhood of Y in W , if any, can be only mild. The fact that the morphism

$$\coprod_{\eta \in \Omega_{(g, n; \beta)}} \Phi_\eta : \coprod_{\eta \in \Omega_{(g, n; \beta)}} \mathfrak{M}(\mathfrak{Y}_0^{rel}, \Gamma_0) \times_{D^r} \mathfrak{M}(\mathfrak{Y}_1^{rel}, \Gamma_1) \longrightarrow \mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; \beta))$$

is natural and surjective ([Li1: Sec. 4] and [Li2: Sec. 4.4]) on the underlying topological space of points with the Zariski topology implies that different choices of W for a given Y give stacks $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; \beta))$ of homeomorphic underlying topologies, (cf. [LMB: Chapter 5]). Since each Φ_η is finite étale of degree a W -independent combinatorial factor $|\text{Eq}(\eta)|$ to its image $\mathfrak{M}(\mathfrak{Y}_0^{rel} \sqcup \mathfrak{Y}_1^{rel}, \eta)$ ([Li1: Sec. 4]), as long as these stacks have the same relative multiplicity to the corresponding image of Φ_η - which is proved to be true and this relative multiplicity of the component of $\mathfrak{M}(\mathfrak{W}_0^{(3)}, (g, n; \beta))$ labelled by η is given by $\mathbf{m}(\eta)$ irrelevant to W ([Li2: Sec. 4.4]) - any detailed difference will not affect the resulting value of Gromov-Witten invariants of Y to be extracted from these stacks for a given $(g, n; \beta)$. (Such indifference is implicit in [Li2] in order to define Gromov-Witten invariants of the degenerate fiber of a family that is family-independent and indeed the degeneration formula implies this indifference as well.)

3 Extracting GW invariants of a conifold from pairs.

The results in Sec. 1 and Sec. 2 together with the constant-under-deformation requirement for any good definition of Gromov-Witten invariants of conifolds implies the following route to extract Gromov-Witten invariants of the conifold Y from the pairs (Y_0, D) and (Y_1, D) , (in this section, by “Gromov-Witten invariants of Y ”, we mean the invariants of Y that would be defined via the intersection theory on the moduli stack $\overline{\mathcal{M}}_{g, n}(Y, \beta)$ of stable maps to Y , should $\overline{\mathcal{M}}_{g, n}(Y, \beta)$ exist and admit a perfect obstruction theory):

$$(g, n; \beta) \Rightarrow \left\{ \left(\mathfrak{M}(\mathfrak{Y}_0^{rel}, \Gamma_0), \mathfrak{M}(\mathfrak{Y}_1^{rel}, \Gamma_1) \right) \right\}_{\eta = (\Gamma_0, \Gamma_1, I) \in \overline{\Omega}_{(g, n; \beta)}} \xrightarrow[\text{from Theorem 2.3}]{\text{gluing formula}} \begin{array}{l} \text{Gromov-Witten} \\ \text{invariants of } Y \\ \text{as would be defined} \\ \text{via } \overline{\mathcal{M}}_{g, n}(Y, \beta). \end{array}$$

There is one last ingredient, though, that we have not yet discussed: when the number n of marked points on the prestable curve is non-zero, the definition of Gromov-Witten invariants of Y involves choices of cycles on Y . In this case, we need to know whether we can choose canonically (perhaps up to some equivalence relation) a set of cycles on $Y_0 \cup_D Y_1$ to feed into the expression in the definition of relative Gromov-Witten invariants for (Y_0, D) and (Y_1, D) in the gluing formula. Let us now turn to this last issue.

Suppose that $Y = W_0$ is in a degeneration $W/\mathbb{A}^1$. Let $\tau_i : A_*(Y_i) \rightarrow A_{*-1}(D)$, $i = 0, 1$, be the group homomorphism defined by taking intersection with D . Let Z be a cycle in W flat over $\mathbb{A}^1$, then $(Z \cdot Y_0) \cdot D = -(Z \cdot Y_1) \cdot D$. Since $Y_0 \cdot D = -Y_1 \cdot D$ on W , the equality $\tau_0(Z \cdot Y_0) = \tau_1(Z \cdot Y_1)$ must hold. This motivates the following definition:

Definition 3.1 [pre-deformable class on $Y_0 \cup_D Y_1$]. Let $\xi_i \in A_*(Y_i)$, $i = 0, 1$. We say that (ξ_0, ξ_1) gives a *pre-deformable class* (i.e. $\xi_0 + \xi_1$) on $Y_0 \cup_D Y_1$ if $\tau_0(\xi_0) = \tau_1(\xi_1)$ in $A_*(D)$. Such (ξ_0, ξ_1) will be called an *admissible pair* of cycle classes with respect to the gluing $Y_0 \cup_D Y_1$.

Readers may note that the image of a pre-deformable map from a (not necessarily connected) prestable curve to $Y_0 \cup_D Y_1$ as defined in [Li1: Sec. 2; in particular, Definition 2.9] gives an example of a pre-deformable class on $Y_0 \cup_D Y_1$.

Recall ([Fu]) that the rational map $\varsigma_0 : Y \dashrightarrow Y_0$ (the inverse of $p_0 : Y_0 \rightarrow Y$) defines a group homomorphism $\varsigma_{0*} : A_*(Y) \rightarrow A_*(Y_0)$ by taking intersection product with the closure of the graph of ς_0 in $Y \times Y_0$.

Definition 3.2 [admissible class on Y]. A class $\xi \in A_*(Y)$ will be called *admissible* if $\varsigma_{0*}(\xi)$ in $A_*(Y_0)$ admits a unique extension to an admissible pair $(\varsigma_{0*}(\xi), \xi')$ with respect to $Y_0 \cup_D Y_1$. Denote the set of all such classes in $A_*(Y)$ by $A_*^{ad}(Y)$.

Note the $\tau_1 : A_*(Y_1) \rightarrow A_*(D)$ is injective on $A_{\geq 1}(Y_1)$ with $\tau_1(A_{\geq 1}(Y_1))$ spanned by $[pt]$, $[(1, 1)\text{-curve}]$, and $[D]$. Consequently, effective classes of constant dimension ξ_1 in $A_{\geq 1}(Y_0)$ whose intersection with D lies in this subspace has a unique extension to an admissible pair (ξ_1, ξ_2) with respect to $Y_0 \cup_D Y_1$. The image of such ξ_1 on Y will span $A_*^{ad}(Y)$. The constant-under-deformation requirement and multi-linearity in the axioms of Gromov-Witten invariants for any variety imply the following:

Corollary 3.3 [extraction of GW invariants of Y from pairs]. *The Gromov-Witten invariants of Y that involve only classes in $A_*^{ad}(Y)$ can be extracted from the relative Gromov-Witten invariants of pair (Y_0, D) and (Y_1, D) by the gluing formula.*

In other words, let $\alpha \in (A_*^{ad}(Y)_{\mathbb{Q}})^{\times n}$ and $\zeta \in A_*(\mathfrak{M}_{g,n})$. Denote by $\Psi_{(g,n;\beta)}^Y(\alpha, \zeta)$ the Gromov-Witten invariant of Y associated to the topological type $(g, n; \beta)$ and classes α and ζ that is defined via *any* standard procedure, i.e. a construction of $\overline{\mathcal{M}}_{g,n}(Y, \beta)$ with a perfect obstruction theory, ..., etc. in such a way that the Gromov-Witten axioms and the constant-under-deformation property are satisfied and that when the construction is applied to smooth variety, it recovers the Gromov-Witten invariants from

the equivalent construction of [B-M] and [L-T]. Let $(\alpha_0, \alpha_1) \in A_*(Y_0)^{\times n} \times A_*(Y_1)^{\times n}$ be the tuple from the admissible pairs with respect to $Y_0 \cup_D Y_1$ of the class in the entries of α . For $\eta = (\Gamma_0, \Gamma_1, I) \in \overline{\Omega}_{(g,n;\beta)}$, assume that $G_\eta^*(\zeta) = \sum_{j \in K_\eta} \zeta_{\eta,0,j} \boxtimes \zeta_{\eta,1,j}$, where $G_\eta : \mathfrak{M}_{\Gamma_0} \times \mathfrak{M}_{\Gamma_1} \rightarrow \mathfrak{M}_{g,n}$ is the natural morphism between the related moduli stack of nodal curves, cf. [Li2: Sec. 0]. Then

$$\Psi_{(g,n;\beta)}^Y(\alpha, \zeta) = \sum_{\eta \in \overline{\Omega}_{(g,n;\beta)}} \frac{\mathbf{m}(\eta)}{|\text{Eq}(\eta)|} \sum_{j \in K_\eta} \left[\Psi_{\Gamma_0}^{Y_0^{rel}}(\alpha_0, \zeta_{\eta,0,j}) \bullet \Psi_{\Gamma_1}^{Y_1^{rel}}(\alpha_1, \zeta_{\eta,1,j}) \right]_0.$$

While the left-hand side of the identity remains to be truly constructed (though one may turn this identity to a *definition* of Gromov-Witten invariants of Y for α admissible if one wishes), the right-hand side is completely determined by and canonically/intrinsically associated to the data: Y , $(g, n; \beta)$, α , and ζ .

Note that when Y is a Calabi-Yau conifold (i.e. it arises from a degeneration of Calabi-Yau 3-folds), the expected dimension of the would-be moduli stack $\overline{\mathcal{M}}_{g,0}(Y, \beta)$ is zero. In this case, no issue of choice of cycles is involved.

Corollary 3.4 [Calabi-Yau conifold]. *When Y is a Calabi-Yau conifold, all the Gromov-Witten invariants of Y associated to the topological type $(g, 0; \beta)$ can be extracted from the relative Gromov-Witten invariants of pairs (Y_0, D) and (Y_1, D) by the gluing formula.*

Remark 3.5 [general conifold]. For a general conifold Y with $\text{Sing}(Y) = \{p_1, \dots\}$, the discussions, expressions, and formulas in Sec. 2 and Sec. 3 generalize immediately by appropriately do the following replacement: D in Y_0 by $\coprod_i D_i$ in Y_0 , (Y_1, D) by isomorphic copies of smooth quadric pairs $(Y_i, D_i) \simeq (Y_1, D)$, summand involving the index 1 of the pair (Y_1, D) by the summation $\sum_i, \dots$, etc., in the expressions involved, where $i = 1, \dots$ corresponds to each conifold singularity p_i of Y .

Remark 3.6 [a general picture]. For a general irreducible singular variety Y , the result in the current work suggests the following picture whose details remain to be studied: an assignment to Y a collection of pairs: a principal pair $(Y_0, D_0 = \cup_i D_{0,i})$, where $Y_0 \rightarrow Y$ is a resolution of Y whose exceptional locus D_0 is a divisor with only simple normal crossing (snc) singularities, and a set of auxiliary smooth (smooth variety)-(snc divisor) pairs (Y_j, D_j) that are related to the germs of the singularities of Y only, and a gluing diagram Γ that encodes the gluing all the these pairs so that (cf. FIGURE 3-1)

- (1) the relative Gromov-Witten invariants of (Y_0, D_0) is obtained from a combination of those for the smooth pairs $(Y_0, D_{0,i})$, $(D_{0,i}, D_{0,i} \cap \cup_{i' \neq i} D_{0,i'})$, $\dots$, etc, and similarly for (Y_j, D_j) (cf. a “quantum inclusion-exclusion principle”);
- (2) the Gromov-Witten invariants of Y is then obtained from a combination of the relative Gromov-Witten invariants of (Y_0, D_0) and (Y_j, D_j) ’s;

- (3) the combination rules in Item (1) and Item (2) above are functorial/universal in the same sense that Jun Li's degeneration formula is functorial/universal (i.e. independent of details of Y away from the singularities, curve classes chosen, cycle classes chosen, ... etc.).




FIGURE 3-1. A gluing construction of Gromov-Witten invariants of a singular variety Y . Since Gromov-Witten invariants arise from A-model topological string theory on the physics side [Wi3], such gluing construction should ring closely with the Atiyah-Segal's formulation of topological quantum field theory ([At] and [Se]; see also [E-G-H] and [B-P]).

References

- [At] M. Atiyah, *Topological quantum field theories*, Publ. Math. I.H.E.S. **68** (1989), pp. 175 - 186.
- [A-GZ-V] V.I. Arnold, S.M. Gusein-Zade, and A.N. Varchenko, *Singularities of differentiable maps*, vol. I. *The classification of critical points, caustics and wave fronts*; vol. II, *Monodromy and asymptotics of integrals*, Birkhäuser, 1988.
- [A-M-V] M. Aganagic, M. Mariño, and C. Vafa, *All loop topological string amplitudes from Chern-Simons theory*, *Commun. Math. Phys.* **247** (2004), pp. 467 - 512.
- [Be] K. Behrend, *Gromov-Witten invariants in algebraic geometry*, *Invent. Math.* **127** (1997), pp. 601 - 617.
- [B-L] V. Braun and C.-H. Liu, *On the isolated singularity of a 7-space obtained by rolling Calabi-Yau 3-folds through extremal transitions*, [hep-th/9801175](#).
- [B-M] K. Behrend and Yu. Manin, *Stacks of stable maps and Gromov-Witten invariants*, *Duke Math. J.* **85** (1996), pp. 1 - 60.
- [B-P] J. Bryan and R. Pandharipande, *The local Gromov-Witten theory of curves*, [math.AG/0411037](#).

- [C-dlO-G-P] P. Candelas, X.C. de la Ossa, P.S. Green, and L. Parkes, *A pair of Calabi-Yau manifolds as an exactly soluble superconformal theory*, Nucl. Phys. **B359** (1991), pp. 21 - 74.
- [C-H] S.D. Cutkosky and J. Herzog, *Cohen-Macaulay coordinate rings of blowup schemes*, Comment. Math. Helv. **72** (1997), pp. 605 - 617.
- [De] O. Debarre, *Higher-dimensional algebraic geometry*, Springer-Verlag, 2001.
- [D-F-G] D.-E. Diaconescu, B. Florea, and A. Grassi, *Geometric transitions and open string instantons*, Adv. Theor. Math. Phys. **6** (2002), pp. 619 - 642.
- [E-G-H] Y. Eliashberg, A. Givental, and H. Hofer, *Introduction to symplectic field theory*, math.SG/0010059.
- [Fu] W. Fulton, *Intersection theory*, Ser. Mod. Surv. Math. 2, Springer-Verlag, 1984.
- [Gr] A. Grothendieck, *Éléments de géométrie algébrique II: Étude globale élémentaire de quelques classes de morphismes*, I.H.E.S. Publ. Math. **8** (1961), pp. 1 - 222.
- [G-H] P. Griffiths and J. Harris, *Principles of algebraic geometry*, John-Wiley & Sons, 1978.
- [G-P] M.B. Green and J. Polchinski, *Summing over world-sheet boundaries*, Phys. Lett. **B335** (1994), pp. 377 - 382.
- [G-R] A. Grassi and M. Rossi, *Large N dualities and transitions in geometry*, math.AG/0209044.
- [G-V] R. Gopakumar and C. Vafa, *On the gauge theory/geometry correspondence*, Adv. Theor. Math. Phys. **3** (1999), pp. 1415 - 1443.
- [Harr] J. Harris, *Algebraic geometry, a first course*, GTM 133, Springer-Verlag, 1992.
- [Hart] R. Hartshorne, *Algebraic geometry*, GTM 52, Springer-Verlag, 1977.
- [Hi] M.W. Hirsch, *Differential topology*, GTM 33, Springer-Verlag, 1976.
- [Ko] J. Kollár, *Flips, flops, minimal models, etc.*, in *Surveys in differential geometry*, **1** (1991), S.-T. Yau and B.H. Lawson, Jr. eds., pp. 113 - 199.
- [K-K-M-SD] G. Kempf, F. Knudsen, D. Mumford, and B. Saint-Donat, *Toroidal embeddings, I*, Lect. Notes Math. 339, Springer-Verlag, 1973.
- [K-M] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts Math. 134, Cambridge Univ. Press, 1998.
- [Li1] J. Li, *Stable morphisms to singular schemes and relative stable morphisms*, J. Diff. Geom. **57** (2001), pp. 509 - 578.
- [Li2] ———, *A degeneration formula of GW-invariants*, J. Diff. Geom. **60** (2002), pp. 199 - 293.
- [L-T] J. Li and G. Tian, *Virtual moduli cycles and Gromov-Witten invariants of algebraic varieties*, J. Amer. Math. Soc. **11** (1998), pp. 119 - 174.
- [L-Y] C.-H. Liu and S.-T. Yau, *A degeneration formula of Gromov-Witten invariants with respect to a curve class for degenerations from blow-ups*, math.AG/0408147.
- [L-MB] G. Laumon and L. Moret-Bailly, *Champs algébriques*, Springer-Verlag, 2000.
- [O-V1] H. Ooguri and C. Vafa, *Knot invariants and topological strings*, Nucl. Phys. **B577** (2000), pp. 419 - 438.
- [O-V2] ———, *World-sheet derivation of a large N duality*, Nucl. Phys. **B641** (2002), pp. 3 - 34.
- [P-S] K.H. Paranjape and V. Srinivas, *Continuous self-maps of quadric hypersurfaces*, in *Proceedings of the Indo-French conference on geometry*, Bombay, 1989, S. Ramanan and A. Beauville eds., pp. 135 - 148, Hindustan Book, Delhi, 1993.
- [Se] G. Segal, *The definition of conformal field theory*, in *Topology, geometry and quantum field theory*, U. Tillmann ed., pp. 421 - 577, (draft widely circulated since 1989), London Math. Soc. Lect. Notes Ser. 308, Cambridge Univ. Press, 2004.
- [Sp] E.H. Spanier, *Algebraic topology*, Springer-Verlag, 1966.

- [St] A. Strominger, *Massless black holes and conifolds in string theory*, *Nucl. Phys.* **B451** (1995), pp. 96 - 108.
- [Va] C. Vafa, *Superstrings and topological strings at large N* , *J. Math. Phys.* **42** (2001), pp. 2798 - 2817.
- [Vi] J.W. Vick, *Homology theory - an introduction to algebraic topology*, Academic Press, 1973.
- [Wi1] E. Witten, *Phases of $N = 2$ theories in two dimensions*, *Nucl. Phys.* **B403** (1993), pp. 159 - 222.
- [Wi2] ———, *Chern-Simons gauge theory as a string theory*, in *The Floer memorial volume*, H. Hofer, C.H. Taubes, A. Weinstein, and Z. Zehnder eds., *Prog. Math.* 133, pp. 637 - 678, Birkhäuser, 1995.
- [Wi3] ———, *Mirror manifolds and topological field theory*, in *Mirror symmetry I*, S.-T. Yau ed., pp. 121 - 160, *Studies Adv. Math.* 9, Amer. Math. Soc. & International Press, 1998.